

Novel type of color solitons with topological charge not coinciding with baryon number and extended chiral group E_χ

Victor Novozhilov and Yuri Novozhilov

V.A.Fock Institute of Physics, St.Petersburg State University, 198504 St.Petersburg, Russia

E-mail: yunovo@pobox.spbu.ru, vnovozhilov@mail.ru

We consider extended chiral group E_χ of gauge plus chiral transformations of the quark path integral in the background gauge and present novel type of color solitons described by the effective action resulting from bosonization. $O(3)$ solitons are formed by a chiral field including parameters with diquark quantum numbers within symplectic subgroup of E_χ , as well as within complete E_χ ; isolated solitons (i.e. solitons in gluonic vacuum) can be stable due to proper asymptotics induced by background vacuum, their topological charge starts from $\pm\frac{4}{3}$ and does not coincide with the baryon number, as if notions of left and right were dependent also on non-abelian directions in such a $SU(2)$ group, where the baryon number defines a diagonal generator.

Understanding world of color solitons is important in a non-perturbative approach to the low energy hadron physics. One of the problems is to include color chiral anomalies [1] in the total (gauge plus chiral, or anomalous) color space and to study changes of gauge space.

In the gauge sector of $SU(2)$ QCD theory, color solitons were recently reported by Faddeev-Niemi [2] and Cho et al. [3]. In the quark chiral sector, the Skyrmion model for constituent quarks (qualitons) was discussed by Kaplan [4], and it was shown recently [5] that isolated color solitons (i.e. solitons on the background of vacuum gluon field) can be classically stabilized by the chromomagnetic vacuum field in the cases of two colors, one flavor and three colors, one flavor, their mass can be evaluated and intersoliton potential

displays confinement behavior.

The aim of this Letter is to present novel type of topological color solitons existing in symplectic subgroup of the extended chiral group $E\chi$, as well as in complete $E\chi$. As a first step we consider isolated solitons which are defined as solitons in vacuum background field. We describe vacuum in a phenomenological way [6] through condensates and assume that cubic gluon condensate is zero.

The extended chiral group $E\chi$ is the group of all gauge and chiral transformations leaving quark lagrangian invariant, if external fields are transformed accordingly [7]. In absence of external fields, $E\chi$ is the group of global color and flavor transformations leaving free quark lagrangian invariant. The basic Dirac spinor Ψ is 8-component, as considered first by Pauli and Gursay : $\Psi = \begin{pmatrix} q \\ \bar{q}^T \end{pmatrix}$, and the Dirac lagrangian L can be rewritten as $L = \frac{1}{2}\Psi^T \hat{F} \Psi$ with $\hat{F}$ represented in the block form $\hat{F} = \begin{pmatrix} C\Phi & -\hat{D}^T \\ \hat{D} & \bar{\Phi}C \end{pmatrix}$, where $\hat{D} = i\gamma^\mu (\partial_\mu + V_\mu + \gamma_5 a_\mu)$ is the Dirac operator with external fields, T means transposition and $\Phi = \gamma^\mu (\phi_{5\mu} + \gamma_5 \phi_\mu)$ contains external vector diquark fields, $\bar{\Phi} = \gamma_0 \Phi^\dagger \gamma_0$.

To avoid difficulties with the Majorana spinors in finding a counterpart of $\hat{F}$ in the Euclidean path integral over Ψ , one should take special care [7] keeping in mind that what is necessary for chiral actions is only to calculate $\det \hat{F}$. To this end we use the hermitean operator $\hat{G} = \begin{pmatrix} \hat{D} & \bar{\Phi} \\ \Phi & \hat{D}_c \end{pmatrix}$, $\hat{D}_c = C^{-1} \hat{D}^T C$, with $\det \hat{G} = \det \hat{F}$ and required positivity properties [7]. Thus, having in mind the chiral anomaly and related effective action, one should study transformations of operator $\hat{G}$ induced by quark transformation $\delta\Psi$

$$\delta\Psi = -\Omega\Psi, \hat{G} \rightarrow \hat{G}' = \exp(-\Xi + \gamma_5\Theta) \hat{G} \exp(\Xi + \gamma_5\Theta) \quad (1)$$

where antihermitean matrices Ω, Ξ, Θ are given by $\Omega = \rho_{11}(\alpha + \gamma_5\chi) + \rho_{22}(\alpha^* - \gamma_5\chi^*) + \rho_{12}(\xi + \gamma_5\omega)C + \rho_{21}(-\xi^* + \gamma_5\omega^*)C$, $\Xi = \rho_{11}\alpha + \rho_{22}\alpha^* + \rho_{12}\xi + \rho_{21}\xi^*$, $\Theta = \rho_{11}\chi - \rho_{22}\chi^* + \rho_{12}\omega - \rho_{21}\omega^*$. We introduced notation: ρ_{ab} is a 2×2 block matrix with elements $(\rho_{ab})_{cd} =$

$\delta_{ac}\delta_{bd}$. We have $\alpha^+ = -\alpha, \chi^+ = -\chi, \xi^T = -\xi, \omega^T = \omega$. The quark baryon number is $\hat{b} = \frac{1}{3}(\rho_{11} - \rho_{22}) = \frac{1}{3}\hat{\rho}_3$.

Transformations with Ξ do not change the quark path integral, while transformations with Θ induce chiral anomalies. The Lie algebras with $\Xi + \gamma_5\Theta$ and $\Xi + \Theta$ are isomorphic. For N_C colors and N_F flavors, generators $\Xi + \Theta$ are in algebra of $U(2N)$, $N = N_C N_F$. Non-anomalous generators α are in algebra of $SU(N)$ and include color $SU(N_C)$; the left-right group $\mathbf{G}_{LR} = SU(N)_L \times SU(N)_R$ has α, χ as generators, while $(\Xi + \Theta)_{sp} = \rho_{11}\alpha + \rho_{22}\alpha^* + \rho_{12}\omega - \rho_{21}\omega^*$ is in algebra of symplectic group $Sp(2N)$, because $(\Xi + \Theta)_{sp}^T i\rho_2 + i\rho_2 (\Xi + \Theta)_{sp} = 0$. In $E\chi$ non-anomalous generators Ξ are in algebra of the orthogonal group $O(2N)$ and include in addition to block diagonal generators α also non-diagonal generators ξ , which arise from non-commutativity of anomalous generators $\Theta_{LR} = \rho_{11}\chi - \rho_{22}\chi^*$ and $\Theta_{sp} = \rho_{12}\omega - \rho_{21}\omega^*$ [7]. Anomalous generators Θ_{sp} of the symplectic group $Sp(2N)$ belong to the coset $Sp(2N)/SU(N)$. We see that there are two distinguished subgroups $\mathbf{G}_{LR}$ and $Sp(2n)$ of $E\chi$ with the same block diagonal non-anomalous generators $\Xi^0 = \rho_{11}\alpha + \rho_{22}\alpha^*$ and different anomalous parts Θ_{LR} and Θ_{sp} , which do not require introduction of additional non-anomalous generators ξ . In the case of entire group $E\chi$, anomalous generators Θ include both χ and ω and belong to the coset $SU(2N)/O(2N)$.

The chiral field is $U = \exp \Theta$; calculation of $\det\{\exp(\gamma_5\Theta)\hat{G}\exp(\gamma_5\Theta)\}/\det \hat{G}$ by integration of anomaly follows the standard bosonization procedure. After eliminating external color axial fields, $a_\mu = 0$, we get the chiral action for $2N$ internal degrees of freedom $W(U) = W_{eff}(U) - W_{wz}$ with an effective lagrangian $L_{eff}(U)$ and the Wess-Zumino term W_{WZ}

$$\begin{aligned}
L_{eff}(U) = & tr_{C,F} \left\{ \frac{f_\omega^2}{4} D_\mu U D^\mu U^{-1} \right. \\
& + \frac{1}{384\pi^2} \left[\frac{1}{2} [UD_\nu U^{-1}, UD_\mu U^{-1}]^2 - (UD_\nu U^{-1} UD^\nu U^{-1})^2 \right] \\
& \left. + \frac{1}{192\pi^2} ([UD^\mu U^{-1}, UD^\nu U^{-1}](\tilde{G}_{\nu\mu} + U\tilde{G}_{\nu\mu}U^{-1}) + \tilde{G}_{\mu\nu}U\tilde{G}^{\mu\nu}U^{-1}) \right\}
\end{aligned} \tag{2}$$

where coefficients contain additional factor $\frac{1}{2}$ coming from square root of quark determinant ; $D_\mu = \partial_\mu + [\tilde{G}_\mu, *]$, $\tilde{G}_\mu = \rho_{11}G_\mu + \rho_{22}(-G_\mu^T) + \rho_{12}\bar{\Phi} + \rho_{21}\Phi$. The kinetic term depends on a phenomenological parameter f_Θ^2 . As in the case of G_{LR} color solitons ..., we do not take into account the term which contains higher derivatives and reflects presence of radial excitations.

Consider the symplectic group $Sp(2N)$ with α in $SU(N)$, $N = N_C N_F$. The chiral field U_{sp} is a mapping from S^3 to $\Theta_{sp} = \rho_+\omega - \rho_-\omega^*$ with $\omega = \omega^T$. For the whole group (with both α and ω) such mappings belong to $\pi_3 = \mathbf{Z}$. Excluding α we get: (a) Two or three colors, one flavor: $\omega = \lambda_a \omega_a$, $a = 1, 3, 4, 6, 8$, no spherical solitons .(b) Two colors, two flavors: $\omega = \Lambda_b \omega_b$, Λ_b are nine traceless symmetrical matrices $\sigma_2 \times \tau_2$ and $\sigma_{\tilde{k}} \times \tau_{\tilde{l}}$; $\tilde{k}, \tilde{l} = 0, 1, 3$; no spherical solitons .(c) Three colors, two flavors; non-anomalous part Ξ_{sp} is given by 12×12 matrix with α in $SU(6)$ built on $SU(3)$ matrices λ_a and flavor Pauli matrices τ_k , while anomalous part Θ_{sp} together with Ξ_{sp} span $Sp(12)$ algebra. Symmetric matrix ω contains fields with diquark quantum numbers associated with both symmetrical matrices $\lambda_a^S \times \tau_k^S$ and both antisymmetrical ones $\lambda_a^A \times \tau_2$. Antisymmetric matrices $\lambda^A = (\lambda_2, -\lambda_5, \lambda_7) \equiv (O_k)$; $(O_k)_{ij} = -i\varepsilon_{kij}$ in color $O(3)$ algebra we combine with unit coordinate vector $r_k, r_k r_k = 1$ into $\hat{r} = O_k r_k$. We retain only those parameters ω , which enter with generators of $O(3)$, introduce the shape function $F_{sp}(R)$ and write the anomalous part Θ_{sp} and the chiral field as

$$\begin{aligned} \Theta_{sp} &= \rho_+ i\tau_2 O_k \vartheta_k - \rho_- i\tau_2 O_k \vartheta_k^* \equiv i\tau_2 \eta \hat{r} F_{sp}, \eta = \rho_1 \cos \chi - \rho_2 \sin \chi \\ U_{sp} &= \exp \Theta_{sp} = 1 + i\tau_2 \eta \hat{r} \sin F + \hat{r}^2 (\cos F - 1), R^2 = x_k x_k \end{aligned} \quad (3)$$

assuming that χ is constant. Isospin matrices $I_k = (\rho_3 \tau_1, \tau_2, \rho_3 \tau_3)$ commute with Θ_{sp} . U_{sp} can describe two conjugate color solitons $U_\pm = \frac{1}{2} (1 \pm \tau_2 \eta) \exp(\pm i\hat{r} F_{sp}(R))$ made out of fields $\vartheta_k, \vartheta_k^*$ with diquark quantum numbers. Θ_{sp} satisfies $\Theta_{sp}^3 = -i\Theta_{sp} F_{sp}^2$. In this special case transformations close in a smaller group, namely $O(6) \sim SU(4)$, and α belong to $U(3) \sim SU(3) \times U(1)$, while Goldstone diquark degrees of freedom $\eta\Theta_{sp}$ belong to the complex

projective space $CP^3 = SU(4)/SU(3) \times U(1)$ [7]. Note that η anticommutes with the quark baryon number $\hat{b} = \frac{1}{3}\rho_3$. Thus, the commutator $[\hat{G}_k, U_{sp}]$ with the background vacuum field $\hat{G}_k = \rho_{11}G_k + \rho_{22}(-G_k^T)$, $G_k^a = V_k\hat{N}^a$, $\hat{N} = \lambda_a N_a$, $N_a N_a = 1$ contains anticommutator $\{N + N^T, \hat{r}\}$ in order to preserve the form of U_{sp} we have to restrict Ξ_{sp} to a common subgroup $O(3)$ of $SU(3) \times SU(3)^*$. This is possible for vacuum background field $G_k^a = V_k\hat{N}^a$, $\hat{N} = \lambda_a N_a$, $tr\hat{N}^3 = 0$ in the gauge $G_0^a = 0$. Then $\hat{G}_k = \rho_{11}G_k + \rho_{22}(-G_k^T) = (\rho_{11} + \rho_{22})V_k\tilde{N}$, where $\tilde{N} = O_l N_l$, $\tilde{N}^3 = \tilde{N}$. Finally, the soliton subgroup $\mathbf{G}_{sol}$ in symplectic $Sp(12)$ is given by $(\Xi + \gamma_5\Theta)_{sp}^{sol} = iO_k(\alpha_k + \gamma_5\tau_2\eta r_k F_{sp}) + i\rho_3\tau_3\alpha_F$, where α_F is a flavor parameter. For comparison, in the left-right group $(\Xi + \gamma_5\Theta)_{LR}$ with both α and χ in $SU(3)$ algebra the analogue soliton subgroup is $(\Xi + \gamma_5\Theta)_{LR}^{sol} = iO_k(\alpha_k + \gamma_5\rho_3 r_k F_{LR})$. Thus, $\mathbf{G}_{sol} = O(3)_L \times O(3)_R \times U(1)_F$, but in the left-right group $\mathbf{G}_{sol}$ the baryon direction ρ_3 is replaced by another one $\tau_2\eta$. If we introduce the baryon $SU(2)_b$ with generators $(\tau_2\rho_1, \tau_2\rho_2, \rho_3)/2i$, then $\tau_2\eta$ can be reached from ρ_3 by rotation.

Complete E_χ group, anomalous generators $\Theta = (\chi, \omega)$. Two or three colors, one flavor: Θ is built on generators $(\rho_3\lambda^A, \lambda^S, \rho_\pm\lambda^S)$, no spherical solitons, no constituent quarks, if generators with $\rho_\pm$ are present and E_χ is not reduced to the left-right subgroup. Two colors, two flavors: no $SU(2)$ and $O(3)$ subgroups in the same sense. Three colors, two flavors: most general Θ is built on generators $(\rho_3\lambda^A\tau^S, \rho_3\lambda^S\tau^A, \rho_\pm\lambda^S\tau^S, \rho_\pm\lambda^A\tau^A)$ and among them there are two $O(3)$ subgroups with generators $\Theta_\pm(O(3)) \sim \frac{1}{2}(1 \pm \zeta)(\lambda_2, -\lambda_5, \lambda_7)$, where $\zeta = \rho_3 1_\tau \cos \kappa + \eta\tau_2 \sin \kappa$, $[\Xi, \zeta] = 0$, and $\eta = \rho_1 \cos \chi - \rho_2 \sin \chi$, $\zeta^2 = 1$, angles κ and χ are two constant parameters; χ comes from the symplectic group $Sp(12)$, while κ describes mixing of left-right and symplectic contributions to ζ : when $\kappa = 0$, no symplectic contribution is present: when $\kappa = \pi/2$, there is no left-right contribution. The chiral field for E_χ in this case is $U = \exp i\zeta\hat{r}F(R)$, it can be obtained by $SU(2)_b$ rotation from U_{sp} or U_{LR} . In this special case $\Xi + \gamma_5\Theta = iO_k(\alpha_k + \gamma_5\zeta r_k F)$, when Ξ is invariant under rotations of baryon $SU(2)_b$, no additional gauge parameters ξ arise from repetition of such transformation. Thus, although we started from different subgroups of the extended chiral group for $N_C = 3, N_F = 2$, the resulting structure of solitonic chiral subgroups is the same:

it is $O(3)_{\tilde{L}} \times O(3)_{\tilde{R}}$, where left $\tilde{L}$ and right $\tilde{R}$ are defined by projectors $\frac{1}{2}(1 \pm \gamma_5 \tilde{\rho})$, where a block matrix $\tilde{\rho}$ with $\tilde{\rho}^2 = 1$, is linear in the Pauli matrices $\rho_1 \tau_2, \rho_2 \tau_2, \rho_3$ of the baryon $SU(2)_b$. The reason is that E_χ includes quarks and antiquarks.

Topological charge $t_\chi(U)$ for a soliton U in the left-right subgroup of E_χ is related to the quark baryon number $b = \rho_3/N_C$

$$t_\chi(U) = \frac{1}{2} \frac{1}{24\pi^2 N_C} \int d^3x \varepsilon_{ijk} \text{tr} \{ \rho_3 U D_i U^+ U D_j U^+ U D_k U^+ \} \quad (4)$$

and coincides with the baryon number of soliton U . For three colors and N_F flavors t_χ starts from $\frac{2}{3}N_F$ (flavor analogous case is skyrmion as dibaryon studied by Balachandran et al.[8]) . The topological charges for solitons in symplectic subgroup $Sp(12)$ and complete group E_χ are winding numbers in directions $\rho_\perp = \tau_2 \eta$ and $\rho_E = \zeta$ within the baryon $SU(2)_b$ -space

$$t_A(U) = \frac{1}{2} \frac{1}{24\pi^2 N_C} \int d^3x \varepsilon_{ijk} \text{tr} \{ \rho_A U \partial_i U^+ U \partial_j U^+ U \partial_k U^+ \}, \quad A = \perp, E. \quad (5)$$

and do not describe solitonic baryon numbers. These winding numbers start from $\left| \frac{4}{3} \right|$. Boundary conditions for the shape functions $F(R)$ in both cases trivially follow from [9].

Finiteness of soliton mass follows from the static effective lagrangian L_{eff} and asymptotic behavior of the shape function $F(R)$ at large R . The mass is given by the positive definite functional, as it can be easily verified; asymptotics of F is defined by the kinetic term. The kinetic term averaged over directions N^k and ν_t of background vacuum field $G_l^k = V_l N^k$, $V_l = -\frac{\pi}{2i} \varepsilon_{ljt} r_j \nu_t R \sqrt{\frac{C_g}{2}}$ in $O(3)$ color and coordinate spaces takes the following form

$$\overline{K} = \frac{1}{4} N_F f_\omega^2 \{ 2F'^2 + 2 \left(\frac{2}{R^2} + \frac{1}{9} \pi^2 C_g R^2 \right) \sin^2 F + \left(\frac{4}{R^2} + \frac{1}{9} \pi^2 C_g R^2 \right) (\cos F - 1)^2 \} \quad (6)$$

where C_g is the gluon condensate. Thus, asymptotic behavior at large R of the shape function $F(R)$ is governed by the same equation as in the case of color $SU(2)$ soliton in [5].

We use the result

$$F \rightarrow (f_0 R)^{-\frac{3}{2}} \exp \left(-\frac{\pi}{3} \sqrt{\frac{C_g}{2}} R^2 \right), R \rightarrow \infty \quad (7)$$

which guarantees that the mass $M = -4\pi \int dR R^2 L_{stat}$ is finite for positive condensate C_g , i.e. for chromomagnetic vacuum field. It follows from the effective lagrangian that the soliton mass is invariant under $SU(2)_b$ rotation.

We have shown that classically stable, finite mass topological solitons exist in the extended chiral group E_χ . In the case of three colors, two flavors their status is described by the special chiral group $O(3)_{\tilde{L}} \times O(3)_{\tilde{R}}$ with notions of left $\tilde{L}$ and right $\tilde{R}$ defined in terms of $\frac{1}{2}(1 \pm \gamma_5 \zeta)$, where ζ is a constant non-abelian charge direction in the $SU(2)_b$ and ρ_3 is baryon number direction. In the case of isolated soliton, $\zeta = \text{const.}$ will be a covariant notion, when ζ commutes with direction $\hat{N}$ of background vacuum $SU(3)$ field, as it is for $\text{tr} \hat{N}^3 = 0$. The vacuum background field defines also an asymptotic behavior of the shape function F ; the field should be chromomagnetic. This pattern based on establishing mapping from S^3 to anomalous part Θ of E_χ and using properties of vacuum background field can be followed in more complicated cases. This expansion of the world of color solitons is not accompanied by the widening of pure QCD gauge space: it is still $SU(3)$; no additional gauge degrees ξ were still required to accomodate novel solitons. However, these degrees ξ are likely to appear, if vacuum is to be described by set of condensates. Novel type of solitons may be essential in discussion of baryon assymetry and baryon number nonconservation.

We thank M.Bordag for the support and hospitality extended to one of the authors (YuN) during his visit to University of Leipzig. We thank M.Bordag and D.Vassilevich for interesting discussions.

[1] S.L.Adler, Phys.Rev. **177**, 2426 (1969); J.S.Bell and R.Jackiw, Nuovo Cim., **A60**, 47 (1969)

- [2] L. Faddeev and A. Niemi, Nature **387**, 58 (1997); R.Battye and P.Sutcliffe, Phys.Rev.Lett.**81**, 4798 (1998); L. Faddeev and A. Niemi, Phys.Rev.Lett. **85**, 3416 (2000); hep-th/0101078
- [3] Y.M. Cho, Phys.Rev.Lett. **87**, 252001 (2001); W.S.Bae, Y.M.Cho, and S.W.Kimm, Phys.Rev. **D65**, 025005 (2002); Y.M.Cho,H.W.Lee and D.G.Pak, Phys.Lett. **B525**, 347 (2002); hep-th/0201179.
- [4] D.B.Kaplan, Phys.Lett. **B235**, 163 (1990); Nucl.Phys. **B351**, 357 (1991); Y.Frishman, A.Hanany and M.Karliner, hep-ph/9507206.
- [5] V.Novozhilov
and Yu.Novozhilov, Phys.Lett. **B522**, 49 (2001), hep-ph/0110006; Teor.Math.Phys. **131**, 62 (2002).
- [6] M.A. Shifman, A.I.Vainshtein and V.I.Zakharov, Nucl.Phys. **B147**,385 (1979)
- [7] Yu.Novozhilov, A. Pronko and D.Vassilevich, Phys.Lett. **B343**, 358 (1995);(E) **B351**, 601 (1995)
- [8] R.Ball, Phys.Lett. **B227** , 445 (1989); Yu. Novozhilov, A. Pronko and D. Vassilevich, Phys.Lett. **B321**, 425 (1994)
- [9] A.P.Balachandran, V.P.Nair, S.G.Rajeev and A.Stern, Phys.Rev. **D27**,1153 (1983);
A.P.Balachandran, F.Lizzi, V.G.J.Rodgers and A.Stern, Nucl.Phys. **B256**, 525 (1985)